

Comparison hexachlorobenzene residues in placentas from Finland and Denmark (1997-2001)

Heqing Shen¹, Katharina Main², Marko Kaleva³, M Ida Schmidt², K Boisen²,
M Chellakooty², Ida Damgaard², H Virtanen³, A-M Haavisto³, Niels Skakkebaek²,
Jorma Toppari³, Karl-Werner Schramm¹

¹GSF-National Research Center for Environment and Health, Institute for Ecological Chemistry

²University Department of Growth & Reproduction, Copenhagen

³Department of Paediatrics and Physiology, University of Turku, Turku

Introduction

Hexachlorobenzene (HCB) continues to be released in to the environment as an industry byproduct, although it has been banned as fungicide in most countries in the 1970s¹. The high persistence and bioaccumulation along the food chain cause that humans are almost exclusively exposed through their diet. Here we present the fetus exposure as the first step to investigate the possible relationship between the male reproductive system toxicity and environmental exposure of this compound. The two-cohort male placenta samples originate from Denmark and Finland, which have different incidences of male urogenital disorders¹.

Material and Methods

The sample collection, preparation, cleanup procedures and HRGC-HRMS analysis conditions have been described elsewhere². HCB was determined in the lipid phase of 112 homogenized whole placenta samples from Turku, Finland² and 168 samples from Copenhagen, Denmark. Statistical analyses and figure plotting were done by Microsoft Excel and Origin 6.0 (*Microsoft, USA*).

Results and Discussion

Statistical analysis shows that the lipid contents for both of Denmark and Finland samples have nearly normal distributions. The geometric means and the average values are the same for the two cohorts respectively. However, the Danish cohort

has slightly lower lipid content than the Finnish one (1.1 and 1.2 g/g 100%). The HCB concentrations for the two cohorts do not show normal distributions. This means that the HCB contents are not apparently fat correlated in the non-fatty placental tissue. Generally, the HCB level in the Danish samples are higher than in the Finnish samples. We calculated this difference on lipid base (ng/g lipid) and wet weight base (10 times of pg/g wet weight) to consider the possible effect of difference in lipid content. However, the average content in the Danish cohort is 1.5 times of that in the Finnish cohort on both lipid base and wet weight base; the difference in the geometric mean of the concentrations is 1.7 -fold with reference to both, lipid and fresh weight. It suggests that HCB exposure level in Danish fetuses is generally higher than in Finland. This geographic difference might suggest geographic difference of HCB sources. It is believed that food is the main source of general exposure³. This difference might reflect the different contaminant levels in diet, similar to the geographic difference of contamination in salmon, which has been described by Hites et al⁴. It is not clear if this difference reflects the higher incidence of male urogenital malformation in Denmark than in Finland. Analysis will be done on the causal relationships of male urogenital malformations with the exposure levels in the further work.

Acknowledgements

This work was supported by the European Commission under the Quality of Life and Management of Living Resources Programme, Key Action 4 (contract number QLK4-CT-2001-00269). We are grateful for the help of Terttu Vartiainen and Hannu Kiviranta (National Public Health Institute, Kuopio, Finland) for organizing the homogenization of the placentas.

References

1. Toppari J, Larsen JC, Christiansen P, Giwercman A, Grandjean P, Guillette Jr LJ, Jegou B, Jensen TK, Jouannet P, Keiding N, Leffers H, McLachlan JA, Meyer O, Müller J, Meyts ER, Scheike T, Sharpe R, Sumpter J & Skakkebaek NE (1996) *Environ Health Perspec* 104, 741-776.
2. Shen H, Main KM, Kaleva M, Virtanen H, Haavisto AM, Skakkebaek NE, Toppari J, Schramm KW (2004) *Placenta* (delivered).
3. Smith AG and Gangolli SD (2002) *Food Chem Toxicol* 40, 767-779
4. Hites RA, Foran JA, Carpenter DO, Hamilton MC, Knuth BA & Schwager SJ (2004) *Science*, 303, 226-229



